

Low-Lying Excitations of Quantum Hall Droplets

J.H.Oaknin¹, L.Martín-Moreno², J.J.Palacios^{1*} and C.Tejedor¹

¹*Departamento de Física de la Materia Condensada, Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain.*

²*Instituto de Ciencia de Materiales (CSIC), Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain.*

(February 1, 2008)

We present the adequate circular representation of charge density excitations in Quantum Hall droplets at filling factor 1. A novel set of operators describes magnetoexcitons with well defined both total and center-of-mass angular momentum. The accuracy of this description is shown by the high overlap of the magnetoexcitons with the exact eigenstates of a quantum dot in a magnetic field obtained from numerical calculations up to 20 electrons. From such magnetoexcitons we get a good understanding of total energies and spectral functions.

PACS number: 73.40.Hm

The electronic structure of quantum Hall droplets (QHD) has been studied extensively [1–14], both due to its relevance in the analysis of quasi-two dimensional (2D) artificial atoms [15,16] and in order to get insight on the behaviour of 2D systems in the quantum Hall regime. Special attention has been given to 2D parabolic quantum dots (PQD) since the non-interacting system is analytically solvable, its high symmetry simplifies the calculations and some of its properties do not depend on the strength of the confinement potential.

In this paper we introduce the set of operators describing low-lying excitations in PQD for filling factor $\nu = 1$. There are two key conditions to obtain these operators: Firstly, the fact that, close to $\nu = 1$, the lowest excitation within each total angular momentum subspace are excitations of integer charge and, secondly, to impose to the eigenstates these operators generate a well defined third component of both total (M) and center-of-mass (M_{CM}) angular momentum. At the end of the paper we discuss several applications of such operators in the analysis and calculation of relevant magnitudes for transport and spectroscopy in QHD.

We consider interacting electrons in a PQD affected by a perpendicular magnetic field B high enough for the N -electron wave functions to comprise only polarised single particle states in the lowest Landau level (LLL). Due to the circular symmetry, it is convenient to work in the symmetric gauge, so that M is a good quantum number. The Hamiltonian can be written as:

$$H = \left[\frac{N}{2} + \frac{1-\gamma}{2}M \right] \hbar \Omega + \sum_{\{m_i\}} \frac{V_{m_1 m_2 m_3 m_4}}{2} c_{m_1}^\dagger c_{m_2}^\dagger c_{m_3} c_{m_4}$$

where the sum runs over the single-particle angular momentum labelling the states of a PQD defined in the xy plane, and characterized by a frequency ω_0 . $\gamma =$

ω_c/Ω , $\omega_c = (eB)/(m^*c)$, $\Omega = (\omega_c^2 + 4\omega_0^2)^{1/2}$, and $l_B = \sqrt{\hbar/(m^*\Omega)}$ is the effective magnetic length. $c^\dagger$ and c are the electron creation and annihilation operators. An additional advantage of PQD is that the electron-electron interaction matrix elements $V_{m_1 m_2 m_3 m_4}$ between states $\{m\}$ in 2D can be computed analytically [17,7]. In this paper we take the interaction as coulombic, unless explicitly stated otherwise. Let us review some of the known properties of this Hamiltonian. Since the interaction only depends on the relative motion of electrons, the QHD spectrum separates in subspaces with different value of M [1,3,6,10]. Due to the reduction of the Hilbert space to the LLL, the dimension of each subspace labelled by (N, M) is finite. Remarkably, wave functions do not depend on ω_0 or B (apart from a rescaling of the length scale), neither does their energy ordering *within* a (N, M) subspace. This property can be used to get information on an infinite system from the results in a PQD. Only the relative positioning of energies for different (N, M) subspaces depend on the values of the parameters defining the QD due to the first term of the Hamiltonian. On increasing B (or, alternatively, decreasing ω_0) the N -electron GS resides in (N, M) subspaces with increasingly larger M [1,3,6,10].

We denote the GS in subspace (N, M) by $|\Phi_M^{(N)}\rangle$. An important fact, which to the best of our knowledge, has never been stated before, is that $\langle N_E \rangle_M = \langle \Phi_M^{(N)} | \sum_{m=-N}^{\infty} c_m^\dagger c_m | \Phi_M^{(N)} \rangle$ is extremely close to an integer. As M increases, $\langle N_E \rangle_M$ increases in steps of height 1, and size typically N , up to a number of steps of the order of $N/2$ beyond which this property fails [18]. We have checked that this property holds not only for the Coulomb interaction, but also for the logarithmic and contact interactions. Therefore, in order to study the low-lying excitations close to M_0 , we can consider a family of variational wave functions, each one of them being a linear combination of Slater determinants with the same value of $\langle N_E \rangle_M$. This additional "symmetry" allows the computation of properties in this system for very large values of N and M .

In this letter we are going to concentrate on the study of the first sector, i.e. when $\langle N_E \rangle_M \approx 1$. Numerically, we find that there is always an eigenstate with the previous property in each subspace (N, M) with $M_0 < M \leq M_0 + N$. Excitations with $\langle N_E \rangle_M = 1$ can be expressed as a representation of the charge density operator projected onto the LLL. They resemble the single-mode excitations previously studied in superfluid ^4He and in the fractional Quantum Hall regime [19]. A set of such operators which

has been put forward [4,5,7,11–13] to describe boson-like edge-wave excitations in PQD is

$$S_{\Delta M}^\dagger = \sum_{k=1}^N (b_k^\dagger/2)^{\Delta M} = \sum_m \sqrt{\frac{(m+\Delta M)!}{m!}} c_{m+\Delta M}^\dagger c_m.$$

$b_k^\dagger = -(x_k + iy_k)/2l_B = -z_k/2$ is, apart from a normalization constant, the ladder operator increasing the single particle angular momentum in the LLL. It has been conjectured [7], and proven for the parabolic electron-electron interaction [11], that these operators represent a bosonization of the low-lying excitations for $\Delta M \ll N$. However the overlap of the states they generate with the true ground state decreases rapidly with ΔM being negligible for $\Delta M \approx N$ (see below). The reason for this is that $S_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$, for $\Delta M > 1$, is not an eigenstate of the third component of the center-of-mass angular momentum. Note that $S_1^\dagger$, when operated on an *exact* eigenstate, gives an eigenstate with M_{CM} increased by 1 [2]. States with well defined M_{CM} should be annihilated by $(S_1)^{M_{CM}}$. It is straightforward to check that the condition $(S_1)^{M_{CM}} S_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle = 0$ is not fulfilled for $\Delta M > 1$.

The previous discussion gives the key idea for finding boson-like excitations (with well defined M and M_{CM}) by means of an operator $J_{\Delta M}^\dagger$ which ejects one electron from the compact state. Since these excitations must have the lowest possible energy, the center-of-mass tends to be at rest, implying the condition $S_1 J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle = 0$. From this condition we get,

$$J_{\Delta M}^\dagger = \sum_m \sqrt{\frac{m!}{(m+\Delta M)!}} c_{m+\Delta M}^\dagger c_m = \sum_{k=1}^N [b_k^\dagger (b_k b_k^\dagger)^{-1}]^{\Delta M}$$

for $1 < \Delta M \leq N$. These operators can be written as $J_{\Delta M}^\dagger = 2^{\Delta M/2} \int dz (z^*)^{-\Delta M} \rho_0(z)$. They are the representation of the projection on the LLL of the density operator $(\rho_0(z))$ on the subset $((z^*)^{-\Delta M})$ of the cylindrical harmonics [20]. From the first quantisation expression for $J_{\Delta M}^\dagger$, we find [21] that it is possible to expand $J_{\Delta M}^\dagger$ in a series which i) for $\Delta M = N \gg 1$ gives the Laughlin [1] wavefunction for a quasi-electron: $| \Phi_{M_0}^{(N)} \rangle = \prod z_i | \Phi_{M_0}^{(N)} \rangle$, and ii) has $S_{\Delta M}^\dagger$ as asymptotic limit for $\Delta M \ll N$. In between these two limits the density charge excitation created by the operator $J_{\Delta M}^\dagger$ consists of a "dressed" electron-hole pair, i.e., a *magnetoexciton*, where the distance between the hole and the electron is proportional to the quantum number ΔM . The electron is always located at the edge of the droplet while the hole moves toward the center of the droplet with ΔM . For $\Delta M = N$, the hole reaches the centre of the droplet and this marks the end of the first sector.

In order to check the quality of $J_{\Delta M}^\dagger$ for describing QHD excitations, table I shows the overlaps $\langle \Phi_M^{(N)} | J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$ and $\langle \Phi_M^{(N)} | S_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$ for $N = 10, 20$ and $\Delta M \leq N$. Note that for $\Delta M \leq 3$, $J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$ is

an exact eigenstate. Using Trugman and Kivelson's [2] procedure to generate eigenstates it is possible to show that this is so for any isotropic pairwise interaction. The conclusion from table I is clear: the branch of lowest excitations are magnetoexcitons with the center-of-mass at rest. $S_{\Delta M}^\dagger$ only describe approximately the excitations in the $\Delta M \ll N$ limit [5,7,11–13] and, as expected, become poor descriptions of the excitations for droplets with small number of electrons. The operators $J_{\Delta M}^\dagger$ represent a bosonization of low-lying excitations of the PQD close to $\nu = 1$. As it happens in other cases of bosonization [22], they do not satisfy bosonic commutation relations strictly; however their expectation values on $| \Phi_{M_0}^{(N)} \rangle$ do.

From the analytical expression for the wavefunctions it is possible to compute their energies $E_M^{(N)} = \langle \Phi_{M_0}^{(N)} | J_{\Delta M} H J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle / \langle \Phi_{M_0}^{(N)} | J_{\Delta M} J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$, for large values of N . Using Wick's theorem, the energy difference between the GS in the (N, M) and (N, M_0) subspaces can be written as:

$$\hbar\Omega(1-\gamma)\Delta M/2 + \epsilon_{d-e}^{(N)}(\Delta M) + \epsilon_{d-h}^{(N)}(\Delta M) + \epsilon_{e-h}^{(N)}(\Delta M).$$

The first term is the single-particle energy difference between the two states. The remaining three terms are the contributions coming from the interaction; they involved combinations of interaction matrix elements, and can be identified as: $\epsilon_{d-e}^{(N)}$ is the interaction between the compact $\nu = 1$ droplet and the ejected electron onto the edge and is practically independent on ΔM . $\epsilon_{d-h}^{(N)}$ is the energy corresponding to the interaction between the $\nu = 1$ droplet and the hole created in it, and decreases almost linearly with ΔM . $\epsilon_{e-h}^{(N)}$ represents the binding energy of the magnetoexciton, and presents a minimum as a function of M . Figure 1 shows the dependence with M of these last three terms for $N = 30$. We find that the total energy difference has the magnetoroton minimum at the same position as $\epsilon_{e-h}^{(N)}$ in the region where the description in terms of $J_{\Delta M}^\dagger$ is valid. The magnetoroton minimum marks the ΔM at which the first instability in the $\nu = 1$ droplet occurs. We find that, for a contact interaction, the position of this intra-Landau magnetoroton at $\nu = 1$ shifts as $2\sqrt{N}$ in the whole range of N .

We consider now another important application of the operators $J_{\Delta M}^\dagger$. Several transport and spectroscopic properties [8,23,24] are controlled, close to $\nu = 1$, by the spectral function

$$A(N, \omega) = \sum_{\eta, m} | \langle \Phi_\eta^{(N)} | c_m^\dagger | \Phi_{M_0}^{(N-1)} \rangle |^2 \delta(\omega - E_\eta^{(N)} + E_0^{(N-1)})$$

where η labels all N -particle states. In spite of the fact that magnetoexcitons $J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$ are not the only excitations in the system, the main contributions to $A(N, \omega)$ come from states of the form $(S_1^\dagger)^j J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle$, that is, from magnetoexcitons and all the states generated from

them by increasing M_{CM} . Figure 2 shows the comparison between the exact spectral function $A(N, \omega)$ (lower panel) and the one calculated including only the preceding states (upper panel), for $N = 8$ and at a value of Ω for which the GS of the 7-electron PQD is the compact state. The identification of the relevant states for the calculation of $A(N, \omega)$ allows i) the analytical calculation of this magnitude for large N and ii) the understanding of the different bunches of peaks spaced in $(\Omega - \omega_c)/2$ and decreasing in intensity that appear in $A(N, \omega)$. After the first high peak coming from the transition between compact states in both $N-1$ and N -particle systems, the first bunch is due to the transition from the compact state of $N-1$ particles to magnetoexcitons of N particles. Each subsequent bunch of peaks in $A(N, \omega)$ is a replica of the preceding one with M_{CM} increased in 1.

Let us now pay attention to the spectrum for large values of ΔM . The lowest-lying states of sectors beyond the first, and even at the end of the first sector ($\Delta M \approx N$) for large values of N , cannot be described in terms of single magnetoexcitons. This is observed in table I where the overlap at $\Delta M = N = 20$ is small [25]. Since an integer number of electrons is ejected from the compact state, one is tempted to look for almost independent bosons in which low-lying states are obtained by successive application of $J_{\Delta M}^\dagger$. Comparing the energies so obtained with the exact ones, the result is satisfactory. However this is not generally the case when comparing wavefunctions. The state $J_{\Delta M_1}^\dagger J_{\Delta M_2}^\dagger |\Phi_{M_0}^{(N)}\rangle$ with $\Delta M_1 + \Delta M_2 > N$ has a small overlap with the state $|\Phi_{M_0 + \Delta M_1 + \Delta M_2}^{(N)}\rangle$ obtained by exact diagonalization. Only when $\Delta M_1 = \Delta M_2 = N$, the overlap is close to 1 and the state can be considered as constituted by two almost non-interacting magnetoexcitons.

Finally, let us discuss one implication of the increasing correlation effects with increasing B . GS changes from the compact state to a magnetoexciton and further to states with several holes in the compact and the same number of electrons out of it. Once spectral functions are known, single electron capacitance experiments in QD [16] can be analyzed. For low temperature, tunneling rates depend only on the spectral weight $\Delta(N) = A(N, \mu(N))$ [14]. This magnitude is shown in fig. 3 as a function of B . For a field in which the GS of both $N-1$ and N electron systems are the compact ones, the spectral weight is obviously 1. For increasing field those two GS melt by ejecting 1 electron. As holes produced in the inner part of the dot are not completely equivalent, $\Delta(N)$ decreases. In some cases the value of the field for which melting appears is significantly different between $N-1$ and N particles so that the spectral weight is drastically quenched. The reduction of $\Delta(N)$ with B explains the experimental quenching of signal amplitude in single electron capacitance experiments for high magnetic fields [16].

In summary, we have found the adequate circular representation of charge density excitations in QHD in

$\nu = 1$. Magnetoexcitons $J_{\Delta M}^\dagger |\Phi_{M_0}^{(N)}\rangle$ turn out to be an excellent description of low-lying excitations. From such wavefunctions it is easy to get energies and spectral functions, i. e. the important information on the electronic structure of QHD.

We thank L. Brey for many useful discussions. This work has been supported in part by the Comisión Interministerial de Ciencia y Tecnología of Spain under contracts No. MAT 91-0905 and MAT 94-0982-C02-01 and by the Commission of the European Communities under contract No. SSC-CT-90-0020.

-
- * Present address: Dep. of Physics, Indiana University, Bloomington, IN 47405.
- [1] R. B. Laughlin, Phys. Rev. Lett. **50**, 231 (1983); Phys. Rev. B **27**, 3383 (1983).
 - [2] S. A. Trugman and S. Kivelson, Phys. Rev. B **31**, 5280 (1985).
 - [3] P. A. Maksym and T. Chakraborty, Phys. Rev. Lett. **65**, 108 (1990); Phys. Rev. B **45**, 1947 (1992).
 - [4] X. G. Wen, Phys. Rev. B **41**, 12838 (1990).
 - [5] M. Stone, Phys. Rev. B **42**, 8399 (1990).
 - [6] U. Merkt *et al.*, Phys. Rev. B **43**, 7320 (1991).
 - [7] M. Stone *et al.*, Phys. Rev. B **45**, 14156 (1992).
 - [8] J. M. Kinaret *et al.*, Phys. Rev. B **46**, 4681 (1992).
 - [9] P. Hawrylak and D. Pfannkuche, Phys. Rev. Lett. **70**, 485 (1993).
 - [10] A. H. MacDonald and M. D. Johnson, Phys. Rev. Lett. **70**, 3107 (1993); *ibid* **71**, 3194 (1993).
 - [11] M. Marsili, Phys. Rev. B **48**, 18051 (1993).
 - [12] A. H. MacDonald *et al.*, Aust. J. Phys. **46**, 581 (1993).
 - [13] S. Giovanazzi *et al.*, Phys. Rev. Lett. **72**, 3230 (1994).
 - [14] J. J. Palacios *et al.*, Phys. Rev. B **50**, 5760 (1994).
 - [15] M. A. Kastner, Rev. Mod. Phys. **64**, 849 (1992).
 - [16] R. C. Ashoori *et al.*, Phys. Rev. Lett. **71**, 613 (1993); Surf. Sc. **305**, 558 (1994).
 - [17] S. M. Girvin and T. Jach, Phys. Rev. B **28**, 4506 (1983).
 - [18] For instance, for $N = 8$, $\langle N_E \rangle_M$ in the first sector is 1 while in the second is $2 > \langle N_E \rangle_M > 1.95$ and $3 > \langle N_E \rangle_M > 2.88$ in the third sector.
 - [19] R. E. Prange and S. M. Girvin, *The Quantum Hall Effect*, (Springer-Verlag, New York, 1987).
 - [20] $S_{\Delta M}^\dagger = 2^{-\Delta M/2} \int dz z^{\Delta M} \rho_0(z)$ are a representation of ρ_0 on other subset ($z^{\Delta M}$) of the cylindrical harmonics.
 - [21] J. H. Oaknin *et al.*, unpublished.
 - [22] G. D. Mahan, "Many particle physics" (Plenum Press, New York, 1981).
 - [23] C. W. J. Beenakker, Phys. Rev. B **44**, 1646 (1991).
 - [24] J. J. Palacios *et al.*, Europhys. Lett. **23**, 495 (1993).
 - [25] An overlap of 0.9982 between the magnetoexciton and the first exact excited state is obtained.

TABLE I. Overlaps between the exact lowest eigenstate $\Phi_M(N)$ and the states obtained applying respectively $J_{\Delta M}^\dagger$ and $S_{\Delta M}^\dagger$ to the compact GS for $N = 10$ and 20 . The small overlap $\langle \Phi_{M_0+20}^{(20)} | J_{20}^\dagger | \Phi_{M_0}^{(20)} \rangle$ is discussed in the text.

ΔM	$N = 10$		$N = 20$	
	$J_{\Delta M}^\dagger$	$S_{\Delta M}^\dagger$	$J_{\Delta M}^\dagger$	$S_{\Delta M}^\dagger$
1	1.000000	1.000000	1.000000	1.000000
2	1.000000	0.994988	1.000000	0.998749
3	1.000000	0.970047	1.000000	0.992503
4	0.999906	0.906678	0.999960	0.976401
5	0.999589	0.788718	0.999752	0.944870
6	0.999285	0.614055	0.999220	0.892297
7	0.999378	0.407942	0.998367	0.814360
8	0.999668	0.219840	0.997480	0.710019
9	0.999831	0.090603	0.992292	0.482037
10	0.999923	0.024995	0.993332	0.315831
11			0.995374	0.196493
12			0.996878	0.114977
13			0.997760	0.082670
14			0.998176	0.057225
15			0.998507	0.037717
16			0.998866	0.023329
17			0.999255	0.013203
18			0.999537	0.006665
19			0.998329	0.003121
20			0.000222	0.000921

FIG. 1. Different contributions to the magnetoexciton energy as a function of ΔM for 30 particles.

FIG. 2. (a) Spectral function for 8 electrons including only magnetoexcitons with rotating center-of-mass $((S_1^\dagger)^j J_{\Delta M}^\dagger | \Phi_{M_0}^{(N)} \rangle)$. (b) Total spectral function including the whole spectrum obtained from the exact diagonalization.

FIG. 3. Many-body contribution to the low-temperature tunneling rates of an electron entering in a QD as given by the spectral weight $\Delta(N)$ as a function of B for different number of electrons.